arXiv:1407.7922v1 [math.AG] 30 Jul 2014

ON THE CANONICAL VOLUME OF 3-FOLDS OF GENERAL TYPE WITH $P_{12} \geq 2$

LINGZI HOU

Abstract. Let V be a nonsingular projective 3-fold of general type. When the pluricanonical section index $\delta(V) > 12$, Chen-Chen [3] has a complete list of the possibility for the weighted basket $\mathbb{B}(V)$. However the possibility of $\mathbb{B}(V)$ could be infinite in the situation $\delta(V) \leq 12$, which is the main challenge to the classification. In this paper we mainly study the case with $\delta(V) = 12$ and show that $\mathrm{Vol}(V) \geq \frac{31}{48048}$, which improves the corresponding result of Chen–Chen [3, Prop. 4.9(5)].

1. Introduction

One of the fundamental problems in birational geometry is to study the distribution of discrete birational invariants of varieties in question. Among those birational invariants, the canonical volume plays the key role. Given a nonsingular projective n-fold Y, the canonical volume

$$\mathrm{Vol}(Y) := \limsup_{n\in\mathbb{Z}^+} \frac{n!\ h^0(Y, mK_Y)}{m^n}.$$

The remarkable theorem, proved by Hacon-McKernan [4], Takayama [6] and Tsuji [7], says that there exists a constant $v_n > 0$ so that $\mathrm{Vol}(Y) \geq v_n$ for any $n \geq 3$. However the constant v_n is not explicitly known except that one has $v_3 \geq \frac{1}{1680}$, which was proved by Chen-Chen [1, 2, 3]. Restricting our interest to 3-folds, we are absorbed by those beautiful calculations of Chen–Chen. Thanks to the guidance of Meng Chen, I am able to consider one of the boundary case here that, when $P_{12} \geq 12$, the possibility for the weighted basket could be infinite and thus it is necessary to slightly extend the method developed by Chen–Chen to find the lower bound of $\mathrm{Vol}(Y)$.

To be precise, we always consider a nonsingular projective 3-fold V of general type. Pick up any minimal model X of V. The so-called weighted basket

$$\mathbb{B}(V) := \mathbb{B}(X) = \{B_X, \chi(\mathcal{O}_X), P_2(X)\}$$

where B_X is the Reid basket (see [5]). Recall from Chen–Chen [3] that the pluricanonical section index

$$\delta(V) = \delta(X) := \min_{m\in\mathbb{Z}^+}\{m | P_m(X) \geq 2\}.$$

The author was supported by National Talents Training Base for Basic Research and Teaching of Natural Science of China, No.J1103105.

By Chen–Chen [1], one has $1 \leq \delta(V) \leq 18$. So far, $\mathbb{B}(V)$ has been completely classified when $\delta(V) > 12$ (see Chen-Chen [3, Theorem 1.4]). When $\delta(V) \leq 12$, though Chen–Chen has given effective lower bounds for K_X^3, it is interesting to see if those are optimal. It is with this motivation that we are going to study the case $\delta(V) = 12$ in this paper. Our main result is the following:

Theorem 1.1. *Let V be a nonsingular projective 3-fold of general type with $P_{12} \geq 2$. Then*

(1) *When $P_{12} = 2$, $\mathbb{B}(V)$ imust be one of the cases listed in 4.3 and 5.3;*
(2) *$Vol(V) \geq \frac{31}{48048}$ and the equality holds if and only if $\mathbb{B}(V) = \{B_H, 4, 0\}$ where*

$$B_H = \{9 \times (1,2), (7,16), (3,7), 2 \times (5,13), 5 \times (1,3), (2,7), (3,11), (1,4)\}.$$

Theorem 1.1 has slightly improved the lower bound $\frac{1}{1560}$ obtained by Chen–Chen [3, Proposition 4.9(5)].

We briefly explain the idea of this paper. Assume $P_{12} \geq 2$. Though $\{\mathbb{B}(V)|\delta(V) = 12\}$ might be an infinite set, we may consider two subcases:(1) $P_{12} \geq 3$; (2) $P_{12} = 2$. The first case is better since Chen–Chen's inequality in [3] can be applied to estimate the lower bound of K_X^3. In the second case, we first deduce that $P_2 = 0$ and $n_{1,6}^0 = 0, 1$. By Chen–Chen's formulae in [1], we can see that $\chi(\mathcal{O}_X)$ is bounded from above. Thus, theoretically, this is a computable problem. In fact, for the situation $n_{1,6}^0 = 0$, we may use Chen-Chen's formula to realize the calculation. When $n_{1,6}^0 = 1$, known formula is not useful for us and we need to deduce the expression for $B^{(n)}$ instead. Fortunately, the result is favorable as there are very few outputs. So we could eventually conclude our main statement.

Essentially affected by Chen–Chen's work, we more or less keep the same notation and concepts as in [1, 2, 3].

Hereby I would like to express my sincere gratitude to professor Meng Chen who guided me to this topic and helped me a lot in completing this paper.

2. **Preliminaries**

We recall the invariants and structure of baskets developed by Chen–Chen in [1].

2.1. **The basket of terminal quotient singularities and packings.**

Definition 2.1. By a 3-dimensional terminal quotient singularity $Q = \frac{1}{r}(1, -1, b)$, we mean one which is analytically isomorphic to the quotient of ($\mathbb{C}^3$, 0) by a cyclic group action ε: $\varepsilon(x, y, z) = (\varepsilon x, \varepsilon^{-1} y, \varepsilon^b z)$, where r is a positive integer, ε is a fixed r-th primitive root of 1, the

integer b is coprime to r and $0 < b < r$. By replacing ε with another primitive root of 1 and changing the ordering of coordinates, we may and do assume that $0 < b \leq \frac{r}{2}$.

Definition 2.2. *A basket B* (of terminal orbifolds) is a collection (allowing weights) of terminal quotient singularities of type $\frac{1}{r_i}(1, -1, b_i)$, $i \in I$ where I is a finite index set. For simplicity, we will always denote a terminal quotient singularity $\frac{1}{r}(1, -1, b)$ simply by the pair (b, r) when no confusion is likely. So we will write a basket as: $B = \{m_i \times (b_i, r_i) | i = 1, 2, \ldots, t\} = \{(b_j, r_j) | j \in J\}, m_i \in \mathbb{Z}^+$, where each m_i denotes the weight. A generalized basket means a collection of pairs of integers (b, r) with $0 < b < r$, but not necessary requiring b, r to be coprime each other.

Definition 2.3. Given a generalized basket (b, r) with $b \leq \frac{r}{2}$ and a fixed integer $n > 0$. Let $\delta := \lfloor \frac{bn}{r} \rfloor$. Then $\frac{\delta+1}{n} > \frac{b}{r} \geq \frac{\delta}{n}$. We define $\Delta^n(b, r) := \delta bn - \frac{(\delta^2+\delta)}{2} r,\ \sigma(B) := \sum_{i \in I} b_i\ ,\ \sigma'(B) := \sum_{i \in I} \frac{b_i^2}{r_i}$.

One can see that $\Delta^n(b, r)$ is a non-negative integer. For a generalized basket $B = \{(b_i, r_i) | i \in I\}$ and a fixed $n > 0$, we define $\Delta^n(B) = \sum_{i \in I} \Delta^n(B_i)$ where $B_i = \{(b_i, r_i)\}$.

Definition 2.4. Given a basket

$$B = \{(b_1, r_1), (b_2, r_2), (b_3, r_3), \ldots, (b_t, r_t)\},$$

we call the basket $B' = \{(b_1 + b_2, r_1 + r_2), (b_3, r_3), \ldots, (b_t, r_t)\}$ *a packing* of B (and B is an unpacking of B'), written as $B \succeq B'$. If, furthermore, $b_1 r_2 - b_2 r_1 = 1$, we call $B \succ B'$ *a prime packing.*

Lemma 2.5. *(Chen–Chen* [1, Lemma 2.8]*) Let $B \succeq B'$ be any packing between generalized baskets. Keep the same notation as above. Then*

(1) $\Delta^n(B) \geq \Delta^n(B')$ *for all* $n \geq 2$*;*
(2) $\Delta^n(B) = \Delta^n(B') \Leftrightarrow \exists\ \delta \geq 0, \delta \in \mathbb{Z}^+$ *so that* $\frac{b_1}{r_1}, \frac{b_2}{r_2} \in [\frac{\delta}{n}, \frac{\delta+1}{n}]$
(3) $\sigma(B) = \sigma(B'); \sigma'(B) = \sigma'(B') + \frac{(r_1 b_2 - r_2 b_1)^2}{r_1 r_2 (r_1 + r_2)}$

Lemma 2.6. *(Chen–Chen* [1, Lemma 2.11]*) Let $B = \{(b_1, r_1), (b_2, r_2)\} \succeq B' = \{(b_1 + b_2, r_1 + r_2)\}$ and $b_1 r_2 - b_2 r_1 = 1$, set $n = r_1 + r_2$. Then $\Delta^n(B) = \Delta^n(B') + 1$.*

Recall that, in Chen–Chen [1, 2.13], an operator $\mathscr{B}^{(n)}(\cdot)$ was defined to obtain the canonical sequence $\{\mathscr{B}^{(n)}(B)\}$ of the given basket B. Roughly speaking, $\mathscr{B}^{(n)}(B)$ is obtained by unpacking B down to the level "n". For $n > 0$, set $B^{(n)} := \mathscr{B}^{(n)}(B)$. So have the canonical sequence:

$$B^{(0)} = \cdots = B^{(4)} \succeq B^{(5)} \succeq \cdots \overset{\alpha_n}{\succeq} B^{(n)} \overset{\alpha_{n+1}}{\succeq} \cdots \succeq B.$$

Define $\epsilon_n(B)$ to be the total number of prime packings belong to the packing process α_n.

Lemma 2.7. *(see Chen–Chen* [1, Claim B in 2.12, Lemma 2.16]*)*

(1) $B^{(n-1)} = \mathscr{B}^{(n-1)}(\mathscr{B}^{(n)}(B)) \succeq \mathscr{B}^{(n)}(B) = B^{(n)}$ *for all* $n \geq 1$.
(2) $\Delta^j(B^{(0)}) = \Delta^j(B)$ *for* $j = 3, 4$.
(3) $\Delta^j(B^{(n-1)}) = \Delta^j(B^{(n)})$ *for* $j < n$
(4) $\Delta^n(B^{(n-1)}) = \Delta^n(B^{(n)}) + \epsilon_n(B)$.
(5) $\Delta^n(B^{(n)}) = \Delta^n(B)$.

2.2. **The weighted basket.** A weighted basket is a formal triple $\mathbb{B} = \{B, \tilde{\chi}, \tilde{\chi}_2\}$ where B is a generalized basket, $\tilde{\chi} \in \mathbb{Z}$ and $\tilde{\chi}_2 \in \mathbb{Z}^+ \cup \{0\}$.

We define the Euler characteristic and K^3 of a formal basket formally as follows. First we define

$$\begin{cases} \chi_2(\mathbb{B}) := \tilde{\chi}_2, \\ \chi_3(\mathbb{B}) := -\sigma(B) + 10\tilde{\chi} + 5\tilde{\chi}_2 \end{cases}$$

and the volume

$$\begin{aligned} K^3(\mathbb{B}) &:= \sigma'(B) - 4\tilde{\chi} - 3\tilde{\chi}_2 + \chi_3(\mathbf{B}) \\ &= -\sigma + \sigma' + 6\tilde{\chi} + 2\tilde{\chi}_2. \end{aligned}$$

For $m \geq 4$, the Euler characteristic $\chi_m(\mathbb{B})$ is defined inductively by

$$\chi_{m+1}(\mathbb{B}) - \chi_m(\mathbb{B}) := \frac{m^2}{2}(K^3(\mathbb{B}) - \sigma'(B)) + \frac{m}{2}\sigma(B) - 2\tilde{\chi} + \Delta^m(B).$$

Clearly, by definition, $\chi_m(\mathbb{B})$ is an integer for all $m \geq 4$ because $K^3(\mathbb{B}) - \sigma'(B) = -4\tilde{\chi} - 3\tilde{\chi}_2 + \chi_3(\mathbb{B})$ and $\sigma = 10\tilde{\chi} + 5\tilde{\chi}_2 - \chi_3(\mathbb{B})$ have the same parity.

Given a $\mathbb{Q}$-factorial terminal 3-fold X, one can associate to X a triple $\mathbb{B}(X) := (B, \tilde{\chi}, \tilde{\chi_2})$ where $B = B(X)$ is Reid's basket, $\tilde{\chi} = \chi(\mathcal{O}_X)$ and $\tilde{\chi_2} = \chi(\mathcal{O}_X(2K_X))$. It's clear that such a triple is a formal basket. The Euler characteristic and K^3 of the formal basket $\mathbb{B}(X)$ are nothing but the Euler characteristic and K^3 of the 3-fold X. For simplicity, we denote $\chi_m(\mathbb{B})$ by $\tilde{\chi}_m$ for all $m \geq 2$. Also denote $K^3(\mathbb{B})$ by $\tilde{K}^3$.

Definition 2.8. Let $\mathbb{B} := (B, \tilde{\chi}, \tilde{\chi_2})$ and $\mathbf{B'} := (B', \tilde{\chi}, \tilde{\chi_2})$ be two formal baskets.

(1) We say that $\mathbf{B'}$ is a packing of $\mathbb{B}$ (written as $\mathbb{B} \succ \mathbf{B'}$) if $B \succ B'$. Clearly "packing" between formal baskets gives a partial ordering.

(2) A formal basket $\mathbb{B}$ is called *positive* if $K^3(\mathbf{B}) > 0$.

(3) A formal basket $\mathbb{B}$ is said to be minimal positive if it is positive and minimal with regard to packing relation.

In the case that X is a minimal projective 3-fold of general type, we see that $K^3(\mathbb{B}(X)) = K_X^3$ and $\tilde{\chi}_m = P_m(X)$ for all $m \geq 2$ by the vanishing theorem.

2.3. **Representation of $B^{(n)}$ in terms of χ and P_m.** Given a minimal projective 3-fold X of general type. Take $\mathbb{B} = \{B(X), \chi, P_2(X)\}$. Then $B^{(0)}$, $B^{(5)}$, $\cdots$, $B^{(12)}$ can be expressed in terms of χ and P_m $(2 \leq m \leq 13)$. The readers are requested to check Chen–Chen [1, P376–P379] for details which are omitted here due to controlling the bulk of this paper.

3. Proof of the main theorem

3.1. **Boundedness.** Let V be a nonsingular projective 3-fold of general type. Pick up any minimal model X of V. Set

$$\mathbb{B}(X) = \{B(X), \chi(\mathcal{O}_X), P_2(X)\}.$$

We would like to prove that $\{\mathbb{B}(X) | P_2(X) = 2\}$ is a finite set.

Lemma 3.1. *If $P_m \leq 1$ for all $m \leq 11$ and $P_{12} = 2$, then $P_2 = 0$.*

Proof. We shall repeatedly use those formulae in [1, P376–P379] and keep the same notations as there.

First of all, we have

$$\epsilon_6 = -3P_2 - P_3 + P_4 + P_5 + P_6 - P_7 - \epsilon = 0.$$

Clearly we have $P_2 \leq 1$. If $P_2 = 1$, then $P_4 = P_5 = P_6 = 1$. It follows that $P_3 = P_7 = \epsilon = 0$. But this is impossible since $P_2 = P_5 = 1$ implies $P_7 \geq 1$. So $P_2 = 0$. □

So we have the following equalities:

$$\begin{cases} \tau := \sigma' - K^3 = 4\chi - P_3 \\ \sigma = 10\chi - P_3 \\ \Delta^3 = 5\chi - 4P_3 + P_4 \\ \Delta^4 = 14\chi - 6P_3 - P_4 + P_5 \\ \Delta^5 = 27\chi - 10P_3 - P_5 + P_6 \\ \Delta^6 = 44\chi - 15P_3 - P_6 + P_7 \\ \Delta^7 = 65\chi - 21P_3 - P_7 + P_8 \\ \Delta^8 = 90\chi - 28P_3 - P_8 + P_9 \\ \Delta^9 = 119\chi - 36P_3 - P_9 + P_{10} \\ \Delta^{10} = 152\chi - 45P_3 - P_{10} + P_{11} \\ \Delta^{11} = 189\chi - 55P_3 - P_{11} + P_{12} \\ \Delta^{12} = 230\chi - 66P_3 - P_{12} + P_{13} \end{cases} \tag{3.1}$$

By Reid's Riemann-Roch formula, if $\chi(\mathcal{O}_X) \leq 0$, then $P_3 \geq 2$ (contradicting to our assumption of Theorem 1.1). By Chen–Chen [2, Corollary3.13], if $\chi(\mathcal{O}_X) = 1$, then $P_{10} \geq 2$. Thus we may always assume $\chi(\mathcal{O}_X) > 1$ from now on.

Lemma 3.2. *If $P_m \leq 1$ for all $m \leq 11$ and $P_{12} = 2$, then $n^0_{1,r} = 0$ for all $r \geq 7$.*

Proof. If there exists an integer $r \geq 7$ such that $n^0_{1,r} > 0$, then $R \geq 6$ and [1, (3.14)] implies:

$$\begin{aligned} & 2P_5 + 3P_6 + P_8 + P_{10} + P_{12} \\ \geq \ & \chi + 4P_3 + P_7 + P_{11} + P_{13} + R. \end{aligned}$$

This implies $P_3 = 0$ and $P_5 = P_6 = 1$. Thus $P_{11} = 1$. Eventually we see $P_8 = P_{10} = 1$ and $P_7 = P_{13} = 0$ But then $P_{13} \geq P_5 \cdot P_8 = 1$ gives a contradiction. So we have $n^0_{1,r} = 0$ for all $r \geq 7$. □

Lemma 3.3. *If $P_m \leq 1$ for all $m \leq 11$, $P_{12} = 2$) and $n^0_{1,6} \neq 0$, then $n^0_{1,5} = 0$ and $n^0_{1,6} = 1$.*

Proof. Clearly the definition of R and the above inequality implies $n^0_{1,6} \leq 1$.

If $n^0_{1,5} \neq 0$, we have

$$R = 2n^0_{1,5} + 5n^0_{1,6} \geq 7.$$

So we get

$$2P_5 + 3P_6 + P_8 + P_{10} \geq P_7 + P_{11} + P_{13} + 7,$$

which says $P_5 = P_6 = P_8 = P_{10} = 1$ and $P_7 = P_{11} = P_{13} = 0$. But then $P_{11} \geq P_5 \cdot P_6 = 1$ (a contradiction). Thus $n^0_{1,5} = 0$. □

Lemma 3.1, Lemma 3.2 and Lemma 3.3 imply the following:

Corollary 3.4. *If $P_m \leq 1$ for all $m \leq 11$ and $P_{12} = 2$, then one of the following occurs:*

(i) *$n^0_{1,r} = 0$ for all $r \geq 6$.*
(ii) *$n^0_{1,5} = 0$, $n^0_{1,6} = 1$ and $n^0_{1,r} = 0$ for $r \geq 7$.*

3.2. **Classification of Case (i).** By the definition of R and ϵ, we have $R = 2n^0_{1,5}$ and $\epsilon = n^0_{1,5}$. To be precise, we have

$$n^0_{1,5} = \epsilon = -P_3 + P_4 + P_5 + P_6 - P_7,$$
$$R = 2\epsilon = -2P_3 + 2P_4 + 2P_5 + 2P_6 - 2P_7.$$

We list all $B^{(n)}$ $(7 \leq n \leq 12)$ here to guide our computation. In explicit, we have

$$\epsilon_7 = \chi - P_3 + P_6 + P_7 - P_8$$

and $B^{(7)} = \{n^7_{b,r} \times (b, r)\}_{\frac{b}{r} \in S^{(7)}}$ has its weights:

$$B^{(7)} \begin{cases} n^7_{1,2} = 2\chi - 3P_3 + 3P_4 - P_5 + P_6 - 2P_7 + P_8 + \eta \\ n^7_{3,7} = \chi - P_3 + P_6 + P_7 - P_8 - \eta \\ n^7_{2,5} = \chi + P_3 - P_4 + P_5 - 3P_6 + P_8 + \eta \\ n^7_{1,3} = 2\chi + 2P_3 - 2P_4 + 2P_6 - P_7 - \eta \\ n^7_{2,7} = \eta \\ n^7_{1,4} = \chi + 2P_3 + P_4 - 2P_5 - P_6 + P_7 - \eta \\ n^7_{1,5} = -P_3 + P_4 + P_5 + P_6 - P_7 \end{cases}$$

where η is the number of prime packings of type $\{(1,3),(1,4)\} \succ \{(2,7)\}$. We have already known

$$\epsilon_8 = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9$$

Thus, taking some prime packings into consideration, $B^{(8)} = \{n^8_{b,r} \times (b,r)\}_{\frac{b}{r} \in S^{(8)}}$ has coefficients:

$$B^{(8)} \begin{cases} n^8_{1,2} = 2\chi - 3P_3 + 2P_4 - P_5 + P_6 - 2P_7 + P_8 + \eta \\ n^8_{3,7} = \chi - P_3 + P_6 + P_7 - P_8 - \eta \\ n^8_{2,5} = \chi + 2P_3 - 4P_6 + P_9 + \eta \\ n^8_{3,8} = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 \\ n^8_{1,3} = 2\chi + 3P_3 - P_4 - P_5 + P_6 - P_7 - P_8 + P_9 - \eta \\ n^8_{2,7} = \eta \\ n^8_{1,4} = \chi + 2P_3 + P_4 - 2P_5 - P_6 + P_7 - \eta \\ n^8_{1,5} = -P_3 + P_4 + P_5 + P_6 - P_7 \end{cases}$$

We know that:

$$\epsilon_9 = -P_3 - P_6 + P_8 + P_9 - P_{10} + \eta$$

Moreover $S^{(9)} - S^{(8)} = \{\frac{4}{9}, \frac{2}{9}\}$. Let ζ be the number of prime packing of type $\{(1,2),(3,7)\} \succ \{(4,9)\}$, then the number of type $\{(1,4),(1,5)\} \succ \{(2,9)\}$ prime packing is $\epsilon_9 - \zeta$. We can get $B^{(9)}$ consisting of the following coefficients:

$$B^{(9)} \begin{cases} n^9_{1,2} = 2\chi - 3P_3 + 2P_4 - P_5 + P_6 - 2P_7 + P_8 + \eta - \zeta \\ n^9_{4,9} = \zeta \\ n^9_{3,7} = \chi - P_3 + P_6 + P_7 - P_8 - \eta - \zeta \\ n^9_{2,5} = \chi + 2P_3 - 4P_6 + P_9 + \eta \\ n^9_{3,8} = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 \\ n^9_{1,3} = 2\chi + 3P_3 - P_4 - P_5 + P_6 - P_7 - P_8 + P_9 - \eta \\ n^9_{2,7} = \eta \\ n^9_{1,4} = \chi + 3P_3 + P_4 - 2P_5 - P_6 + P_7 - P_8 - P_9 + P_{10} - 2\eta + \zeta \\ n^9_{2,9} = -P_3 - P_6 + P_8 + P_9 - P_{10} + \eta - \zeta \\ n^9_{1,5} = P_4 + P_5 + 2P_6 - P_7 - P_8 - P_9 + P_{10} - \eta + \zeta \end{cases}$$

One has

$$\epsilon_{10} = -P_4 - P_5 + P_6 + P_7 + P_{10} - P_{11} - \eta$$

and then $B^{(10)}$ consists of the following coefficients:

$$B^{(10)}\begin{cases} n^{10}_{1,2} = 2\chi - 3P_3 + 2P_4 - P_5 + P_6 - 2P_7 + P_8 + \eta - \zeta \\ n^{10}_{4,9} = \zeta \\ n^{10}_{3,7} = \chi - P_3 + P_6 + P_7 - P_8 - \eta - \zeta \\ n^{10}_{2,5} = \chi + 2P_3 - 4P_6 + P_9 + \eta \\ n^{10}_{3,8} = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 \\ n^{10}_{1,3} = 2\chi + 3P_3 - 2P_7 - P_8 + P_9 - P_{10} + P_{11} - \eta \\ n^{10}_{3,10} = -P_4 - P_5 + P_6 + P_7 + P_{10} - P_{11} - \eta \\ n^{10}_{2,7} = P_4 + P_5 - P_6 - P_7 - P_{10} + P_{11} + 2\eta \\ n^{10}_{1,4} = \chi + 3P_3 + P_4 - 2P_5 + P_7 - P_8 - P_9 + P_{10} - 2\eta + \zeta \\ n^{10}_{2,9} = -P_3 - P_6 + P_8 + P_9 - P_{10} + \eta - \zeta \\ n^{10}_{1,5} = P_4 + P_5 + 2P_6 - P_7 - P_8 - P_9 + P_{10} - \eta + \zeta \end{cases}$$

By computing $\Delta^{11}(B^{(10)})$, I get

$$\epsilon_{11} = \chi - P_5 - P_6 + P_9 + P_{11} - P_{12} - \zeta$$

Let α be the number of prime packing of type $\{(1,2),(4,9)\} \succ \{(5,11)\}$, and β be the number of prime packing of type $\{(1,3),(3,8)\} \succ \{(4,11)\}$. Then, I get $B^{(11)}$ with

$$B^{(11)}\begin{cases} n^{11}_{1,2} = 2\chi - 3P_3 + 2P_4 - P_5 + P_6 - 2P_7 + P_8 + \eta - \zeta - \alpha \\ n^{11}_{5,11} = \alpha \\ n^{11}_{4,9} = \zeta - \alpha \\ n^{11}_{3,7} = \chi - P_3 + P_6 + P_7 - P_8 - \eta - \zeta \\ n^{11}_{2,5} = \chi + 2P_3 - 4P_6 + P_9 + \eta \\ n^{11}_{3,8} = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 - \beta \\ n^{11}_{4,11} = \beta \\ n^{11}_{1,3} = 2\chi + 3P_3 - 2P_7 - P_8 + P_9 - P_{10} + P_{11} - \eta - \beta \\ n^{11}_{3,10} = -P_4 - P_5 + P_6 + P_7 + P_{10} - P_{11} - \eta \\ n^{11}_{2,7} = -\chi + P_4 + 2P_5 - P_7 - P_9 - P_{10} + P_{12} + 2\eta + \zeta + \alpha + \beta \\ n^{11}_{3,11} = \chi - P_5 - P_6 + P_9 + P_{11} - P_{12} - \zeta - \alpha - \beta \\ n^{11}_{1,4} = 3P_3 + P_4 - P_5 + P_6 + P_7 - P_8 - 2P_9 + P_{10} - P_{11} + P_{12} - 2\eta + 2\zeta + \alpha + \beta \\ n^{11}_{2,9} = -P_3 - P_6 + P_8 + P_9 - P_{10} + \eta - \zeta \\ n^{11}_{1,5} = P_4 + P_5 + 2P_6 - P_7 - P_8 - P_9 + P_{10} - \eta + \zeta \end{cases}$$

Finally, since

$$\epsilon_{12} = -\chi - 2P_3 - P_4 + P_5 + P_8 + P_{12} - P_{13} + \eta$$

$$B^{(12)}\begin{cases} n^{12}_{1,2} = 2\chi - 3P_3 + 2P_4 - P_5 + P_6 - 2P_7 + P_8 + \eta - \zeta - \alpha \\ n^{12}_{5,11} = \alpha \\ n^{12}_{4,9} = \zeta - \alpha \\ n^{12}_{3,7} = 2\chi + P_3 + P_4 - P_5 + P_6 + P_7 - 2P_8 - P_{12} + P_{13} - 2\eta - \zeta \\ n^{12}_{5,12} = -\chi - 2P_3 - P_4 + P_5 + P_8 + P_{12} - P_{13} + \eta \\ n^{12}_{2,5} = 2\chi + 4P_3 + P_4 - P_5 - 4P_6 - P_8 + P_9 - P_{12} + P_{13} \\ n^{12}_{3,8} = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 - \beta \\ n^{12}_{4,11} = \beta \\ n^{12}_{1,3} = 2\chi + 3P_3 - 2P_7 - P_8 + P_9 - P_{10} + P_{11} - \eta - \beta \\ n^{12}_{3,10} = -P_4 - P_5 + P_6 + P_7 + P_{10} - P_{11} - \eta \\ n^{12}_{2,7} = -\chi + P_4 + 2P_5 - P_7 - P_9 - P_{10} + P_{12} + 2\eta + \zeta + \alpha + \beta \\ n^{12}_{3,11} = \chi - P_5 - P_6 + P_9 + P_{11} - P_{12} - \zeta - \alpha - \beta \\ n^{12}_{1,4} = 3P_3 + P_4 - P_5 + P_6 + P_7 - P_8 - 2P_9 + P_{10} - P_{11} + P_{12} - 2\eta + 2\zeta + \alpha + \beta \\ n^{12}_{2,9} = -P_3 - P_6 + P_8 + P_9 - P_{10} + \eta - \zeta \\ n^{12}_{1,5} = P_4 + P_5 + 2P_6 - P_7 - P_8 - P_9 + P_{10} - \eta + \zeta \end{cases}$$

We now explain the effectivity of the possible computation. We have $\epsilon_{10} + \epsilon_{12} \geq 0$ which gives rise to:

$$2P_5 + 3P_6 + P_8 + P_{10} + 2 \geq \chi + 4P_3 + P_7 + P_{11} + P_{13} + R \tag{3.2}$$

Inserting the expression of R, one has:

$$\chi \leq -2P_3 - 2P_4 + P_6 + P_7 + P_8 + P_{10} - P_{11} - P_{13} + 2.$$

If the equality holds, one has $P_3 = P_4 = P_{11} = P_{13} = 0$ and $P_6 = P_7 = P_8 = P_{10} = 1$. But then $P_{13} \geq P_6 \cdot P_7 = 1$, a contradiction. Thus we have

$$\chi \leq 5 \tag{3.3}$$

Considering $n^7_{1,4} \geq 0$, we get:

$$\eta \leq \chi + 2P_3 + P_4 - 2P_5 - P_6 + P_7.$$

So,

$$\eta \leq \chi + 4 \tag{3.4}$$

We have $n^7_{2,9} \geq 0$, which implies:

$$\zeta \leq \chi + P_3 + P_4 - 2P_5 - 2P_6 + P_7 + P_8 + P_9 - P_{10}.$$

Then,

$$\zeta \leq \chi + 4. \tag{3.5}$$

Similarly, by $n^{11}_{4,9} \geq 0$ and $n^{11}_{3,8} \geq 0$, we get:

$$\alpha \leq \zeta \tag{3.6}$$

$$\beta \leq 3 \tag{3.7}$$

Finally we have:

$$P_{13} \leq -\chi - 2P_3 - 2P_4 + P_6 + P_7 + P_8 + P_{10} - P_{11} + 2 \leq -\chi + 6. \tag{3.8}$$

Inequalities (3.3) $\sim$ (3.8) tell us that $\{B^{(12)}\}$ is a computable finite set. Our computer program using Matlab outputs totally 149 classes as follows. We also list all possible minimal positive baskets dominated by these 149 classes of $B^{(12)}$. The last volume provides their volumes.

No	$(\chi, P_3, P_4, \ldots, P_{11})$	$B^{(12)} = (n^{12}_{1,2}, n^{12}_{5,11}, \ldots, n^{12}_{1,5})$	K^3
1	2,0,0,0,0,0,0,0,0,0	4,0,0,2,0,2,0,0,4,0,0,0,2,0,0	1/210
2	2,0,1,0,0,0,1,0,1,0	7,0,0,1,0,2,0,0,2,0,0,0,3,0,1	1/420
3	2,0,0,1,0,0,0,1,1,0	3,0,0,1,1,2,0,0,4,0,0,0,0,0,1	1/420
a		(3,7),(5,12)$\succ$(8,19)	1/570
b		(5,12),2(2,5)$\succ$(9,22)	1/1155
4	2,0,0,1,0,0,0,1,1,0	3,0,0,2,0,3,0,0,4,0,0,0,0,0,1	1/210
5	2,0,0,1,0,1,1,0,1,1	2,0,0,1,1,1,2,0,1,0,0,0,1,0,0	1/420
a		(3,7),(5,12)$\succ$(8,19)	1/570
b		(5,12),(2,5)$\succ$(7,17)	1/714
6	2,0,0,1,0,1,1,0,1,1	2,0,0,2,0,2,2,0,1,0,0,0,1,0,0	1/210
a		2(2,5),2(3,8)$\succ$2(5,13)	1/1092
7	2,0,0,0,0,0,1,0,1,1	5,0,0,0,1,1,0,1,2,0,0,0,2,0,0	1/220
a		(5,12),(2,5)$\succ$(7,17);(4,11),(1,3)$\succ$(5,14)	1/714
b		(4,11),2(1,3)$\succ$(6,17)	1/1020
8 ,	2,0,0,0,0,0,1,0,1,1	5,0,0,1,0,2,0,1,2,0,0,0,2,0,0	8/1155
a		(4,11),2(1,3)$\succ$(6,17)	2/595
9	2,0,0,1,0,1,1,1,1,1	1,0,1,0,1,2,1,0,2,0,0,0,1,0,0	1/360
a		(5,12),2(2,5)$\succ$(9,22)	1/792
b		(2,5),(3,8)$\succ$(5,13)	1/1170
10	2,0,0,1,0,1,1,1,1,1	1,0,1,1,0,3,1,0,2,0,0,0,1,0,0	13/2520
a		(4,9),(3,7)$\succ$(7,16);(6,15),(3,8)$\succ$(9,23)	1/1104
11	2,0,0,0,0,0,1,1,1,1	3,1,0,0,0,3,0,0,4,0,0,0,2,0,0	1/165
a		(1,2),(5,11)$\succ$(6,13)	1/390
12	2,0,0,0,0,0,0,0,1,0	5,0,0,0,1,2,0,0,3,0,1,0,1,0,0	1/210
a		(5,12),2(2,5)$\succ$(9,22)	1/308
13	2,0,0,0,0,0,0,0,1,0	5,0,0,1,0,3,0,0,3,0,1,0,1,0,0	1/140
14	2,0,0,0,1,0,0,1,1,0	6,0,0,2,0,0,0,0,4,1,0,0,0,0,1	1/210
a		(1,3),(3,10)$\succ$(4,13)	1/455
15	2,0,1,0,1,0,1,1,1,0	9,0,0,1,0,0,0,0,3,0,1,0,0,1,1	1/630
16	2,0,1,0,1,1,1,1,1,1	7,0,0,2,0,0,0,0,2,0,0,1,0,1,0	1/1386
17	2,0,0,0,1,1,0,1,0,1	3,0,1,2,0,0,0,0,4,0,1,0,1,0,0	1/252
a		(4,9),2(3,7)$\succ$(10,23)	5/1932
18	2,0,1,0,1,1,1,1,1,1	6,0,1,1,0,0,0,0,2,0,1,0,2,0,1	1/630
19	2,0,0,0,1,0,1,1,1,1	6,0,1,0,0,0,1,0,4,0,1,0,0,0,1	19/2520
20	2,0,0,0,1,1,1,1,1,1	4,0,1,1,0,0,1,0,2,1,0,0,1,0,0	13/2520
a		(4,9),(3,7)$\succ$(7,16);(1,3),(3,10)$\succ$(4,13)	1/624
b		2(1,3),(3,10)$\succ$(5,16)	1/1008
21	2,0,0,0,1,1,0,0,1,1	5,0,0,2,0,0,1,0,2,0,2,0,0,0,0	1/168
22	2,0,0,0,1,1,0,1,1,1	4,0,1,0,1,0,0,0,3,0,2,0,0,0,0	1/252
23	2,0,0,0,1,1,0,1,1,1	4,0,1,1,0,1,0,0,3,0,2,0,0,0,0	2/315
a		(4,9),(3,7)$\succ$(7,16)	3/560
24	3,0,0,1,0,0,1,0,1,0	6,0,0,2,0,3,2,0,4,0,0,0,1,0,1	1/210
a		(4,10),2(3,8)$\succ$(10,26)	1/1092
25	3,0,0,1,0,0,1,1,1,0	6,0,0,2,0,4,0,1,4,0,0,0,0,1,0	13/3465
a		(4,11),(1,3)$\succ$(5,14)	1/630
26	3,0,0,1,0,1,1,0,1,1	4,0,0,3,0,3,1,1,2,0,0,0,2,0,0	31/9240
a		(2,5),(3,8)$\succ$(5,13)	43/30030
b		(4,11),(1,3)$\succ$(5,14)	1/840
27	3,0,0,1,1,0,1,1,1,1	7,0,0,3,0,0,1,1,5,0,0,0,0,0,2	31/9240

a		(3,8),(4,11)≻(7,19)	11/3990
b		(4,11),(1,3)≻(5,14)	1/840
28	3,0,0,0,0,0,1,0,0,0	6,0,1,1,0,3,1,0,5,0,0,0,3,0,0	13/2520
a		(4,9),(3,7)≻(7,16);(6,15),(3,8)≻(9,23)	1/1104
29	3,0,0,1,0,0,1,1,1,0	5,0,1,1,0,4,1,0,5,0,0,0,1,0,1	13/2520
a		(4,9),(3,7)≻(7,16);(6,15),(3,8)≻(9,23)	1/1104
b		(8,20),(3,8)≻(11,28)	1/630
30	3,0,0,0,1,0,1,1,0,1	7,0,1,2,0,0,0,1,6,0,0,0,2,0,1	13/3465
a		(4,9),(6,14)≻(10,23)	3/1265
b		(4,11),(1,3)≻(5,14)	1/630
31	3,0,0,1,0,1,1,1,1,1	3,0,1,2,0,4,0,1,3,0,0,0,2,0,0	13/3465
a		(4,9),(6,14)≻(10,23)	3/1265
b		(4,11),(1,3)≻(5,14)	1/630
32	3,0,0,0,1,0,1,1,0,1	6,1,0,2,0,0,1,0,7,0,0,0,2,0,1	23/9240
33	3,0,0,1,0,1,1,1,1,1	2,1,0,2,0,4,1,0,4,0,0,0,2,0,0	23/9240
34	3,0,0,0,0,0,1,1,0,0	5,0,2,0,0,4,0,0,6,0,0,0,3,0,0	1/180
35	3,0,0,0,1,0,0,0,0,0	8,0,0,3,0,0,1,0,6,0,1,0,1,0,1	1/280
36	3,0,0,0,0,0,0,0,1,0	7,0,0,2,0,4,0,0,5,0,0,1,1,0,0	29/4620
a		(3,11),(1,4)≻(4,15)	1/210
37	3,0,0,1,0,1,0,0,1,0	4,0,0,2,1,3,1,0,3,0,1,0,1,0,0	1/840
38	3,0,1,0,1,0,1,0,1,0	11,0,0,2,0,0,1,0,4,0,1,0,2,0,2	1/840
39	3,0,0,0,1,0,1,0,0,0	9,0,0,2,0,0,2,0,5,0,1,0,0,1,0	1/252
40	3,0,0,1,0,1,0,0,1,0	4,0,0,3,0,4,1,0,3,0,1,0,1,0,0	1/280
a		(2,5),(3,8)≻(5,13)	3/1820
41	3,0,0,0,1,0,1,0,1,0	9,0,0,2,0,0,2,0,4,1,0,0,1,0,1	1/210
a		(1,3),(3,10)≻(4,13)	1/455
42	3,0,0,0,1,0,1,0,1,1	9,0,0,2,0,0,2,0,5,0,0,1,0,0,1	29/4620
43	3,0,0,1,1,1,1,0,1,1	6,0,0,3,0,0,3,0,3,0,1,0,0,0,1	1/280
44	3,0,0,0,1,0,1,1,1,0	9,0,0,1,1,0,0,1,4,1,0,0,0,1,0	19/13860
a		(3,7),(5,12)≻(8,19)	7/9405
45	3,0,0,0,1,0,1,1,1,0	9,0,0,2,0,1,0,1,4,1,0,0,0,1,0	13/3465
a		(4,11),(1,3)≻(5,14)	1/630
b		(1,3),(3,10)≻(4,13)	27/22733
46	3,0,0,0,1,0,1,0,1,1	9,0,0,2,0,0,1,1,4,0,1,0,1,0,1	53/9240
a		(3,8),(4,11),(1,3)≻(8,22)	3/1540
b		(4,11),3(1,3)≻(7,20)	1/840
47	3,0,0,0,1,1,1,0,1,1	7,0,0,3,0,0,1,1,2,1,0,0,2,0,0	31/9240
a		(3,8),(4,11)≻(7,19)	11/3990
b		(4,11),(1,3)≻(5,14)	1/840
c		(1,3),(3,10)≻(4,13)	19/24024
48	3,0,0,1,1,1,0,1,1,1	5,0,0,4,0,1,0,1,4,0,1,0,0,0,1	1/462
49	3,0,0,0,1,0,0,1,0,0	7,0,1,2,0,1,0,0,7,0,1,0,1,0,1	1/252
a		(4,9),2(3,7)≻(10,23)	5/1932
50	3,0,0,0,1,0,1,1,0,0	8,0,1,0,1,0,1,0,6,0,1,0,0,1,0	1/504
51	3,0,0,1,0,1,0,1,1,0	3,0,1,1,1,4,0,0,4,0,1,0,1,0,0	1/630
a		(3,7),(5,12)≻(8,19)	23/23940
52	3,0,0,0,1,0,1,1,1,0	8,0,1,0,1,0,1,0,5,1,0,0,1,0,1	1/360
53	3,0,0,0,1,0,1,1,1,1	8,0,1,0,1,0,1,0,6,0,0,1,0,0,1	17/3960
54	3,0,0,0,0,0,1,0,1,0	7,0,1,0,0,4,1,0,4,0,1,0,2,0,0	19/2520
a		4(2,5),(3,8)≻(11,28)	1/252
55	3,0,0,0,1,0,1,1,0,0	8,0,1,1,0,1,1,0,6,0,1,0,0,1,0	11/2520
a		(4,9),(3,7)≻(7,16);(2,5),(3,8)≻(5,13)	19/13104
56	3,0,0,1,0,1,0,1,1,0	3,0,1,2,0,5,0,0,4,0,1,0,1,0,0	1/252
a		(4,9),2(3,7)≻(10,23)	5/1932
57	3,0,0,0,1,1,1,0,0,1	6,0,1,2,0,0,2,0,4,0,1,0,2,0,0	1/252
a		(4,9),2(3,7)≻(10,23)	5/1932
58	3,0,0,0,1,0,1,1,1,0	8,0,1,1,0,1,1,0,5,1,0,0,1,0,1	13/2520

a		(4,9),(3,7)≻(7,16);(2,5),(3,8)≻(5,13)	7/3120
b		2(1,3),(3,10)≻(5,16)	1/1008
59	3,0,0,0,1,0,1,1,1,1	8,0,1,1,0,1,1,0,6,0,0,1,0,0,1	37/5544
a		(4,9),(3,7)≻(7,16);(2,5),(3,8)≻(5,13)	43/11440
60	3,0,0,1,1,1,1,1,1,1	5,0,1,1,1,0,2,0,4,0,1,0,0,0,1	1/630
a		(3,7),(5,12)≻(8,19)	23/23940
61	3,0,0,1,1,1,1,1,1,1	5,0,1,2,0,1,2,0,4,0,1,0,0,0,1	1/252
a		(4,9),2(3,7)≻(10,23)	5/1932
b		(2,5),2(3,8)≻(8,21)	1/630
62	3,0,0,0,1,0,1,1,1,1	8,0,1,0,1,0,0,1,5,0,1,0,1,0,1	13/3465
a		(4,11),(1,3)≻(5,14)	1/630
63	3,0,0,0,1,0,1,1,1,1	8,0,1,1,0,1,0,1,5,0,1,0,1,0,1	17/2772
a		(4,9),(3,7)≻(7,16);(4,11),2(1,3)≻(6,17)	3/1904
b		(4,11),4(1,3)≻(8,23)	5/5796
64	3,0,0,0,1,1,1,1,1,1	6,0,1,2,0,1,0,1,3,1,0,0,2,0,0	13/3465
a		(4,9),2(3,7)≻(10,23)	3/1265
b ,		(4,11),(1,3)≻(5,14)	1/630
c		(1,3),(3,10)≻(4,13)	27/22733
65	3,0,0,0,1,0,1,1,1,1	7,1,0,0,1,0,1,0,6,0,1,0,1,0,1	23/9240
66	3,0,0,0,1,0,1,1,1,1	7,1,0,1,0,1,1,0,6,0,1,0,1,0,1	3/616
a		(2,5),(3,8)≻(5,13)	59/20020
67	3,0,0,0,1,1,1,1,1,1	5,1,0,2,0,1,1,0,4,1,0,0,2,0,0	23/9240
68	3,0,0,0,1,1,1,1,0,1	5,0,2,1,0,1,1,0,5,0,1,0,2,0,0	11/2520
a		2(4,9),(3,7)≻(11,25);(2,5),(3,8)≻(5,13)	8/6825
69	3,0,0,0,1,0,0,0,1,0	9,0,0,1,1,0,1,0,5,0,2,0,0,0,1	1/280
a		(3,7),(5,12)≻(8,19)	47/15960
70	3,0,0,0,1,0,0,0,1,0	9,0,0,2,0,1,1,0,5,0,2,0,0,0,1	1/168
a		(2,5),(3,8)≻(5,13)	11/2730
71	3,0,0,0,1,1,0,0,1,0	7,0,0,3,0,1,1,0,3,1,1,0,1,0,0	1/280
a		(2,5),(3,8)≻(5,13);(3,10),(2,7)≻(5,17)	5/6188
b		(1,3),(3,10)≻(4,13)	11/10920
72	3,0,0,0,1,1,0,0,1,1	7,0,0,3,0,1,1,0,4,0,1,1,0,0,0	47/9240
a		(2,5),(3,8)≻(5,13);(2,7),(3,11)≻(5,18)	2/819
73	3,0,0,0,1,1,0,0,1,1	7,0,0,3,0,1,0,1,3,0,2,0,1,0,0	1/220
		(4,11),2(1,3)≻(6,17)	1/1020
74	3,0,0,0,1,0,0,1,1,0	8,0,1,0,1,1,0,0,6,0,2,0,0,0,1	1/252
a		(5,12),(2,5)≻(7,17)	16/5355
75	3,0,0,0,1,0,0,1,1,0	8,0,1,1,0,2,0,0,6,0,2,0,0,0,1	2/315
a		(4,9),(3,7)≻(7,16)	3/560
76	3,0,0,0,1,1,0,1,1,0	6,0,1,2,0,2,0,0,4,1,1,0,1,0,0	1/252
a		(4,9),2(3,7)≻(10,23);(3,10),(2,7)≻(5,17)	41/23460
b		(1,3),(3,10)≻(4,13)	23/16380
77	3,0,0,0,1,1,1,0,1,1	7,0,1,0,1,0,2,0,3,0,2,0,1,0,0	1/252
78	3,0,0,0,1,1,0,1,1,1	6,0,1,2,0,2,0,0,5,0,1,1,0,0,0	19/3465
a		(4,9),2(3,7)≻(10,23);(2,7),(3,11)≻(5,18)	7/2070
79	3,0,0,0,1,1,1,0,1,1	7,0,1,1,0,1,2,0,3,0,2,0,1,0,0	2/315
a		(4,9),(3,7)≻(7,16);(2,5),2(3,8)≻(8,21)	1/336
80	3,0,0,0,1,1,0,1,1,1	5,1,0,2,0,2,0,0,5,0,2,0,1,0,0	17/4620
81	3,0,0,0,1,1,1,1,1,1	6,0,2,0,0,2,1,0,4,0,2,0,1,0,0	17/2520
a		2(2,5),(3,8)≻(7,18)	1/252
82	4,0,0,0,1,0,1,0,0,0	11,0,0,3,0,1,2,0,7,0,0,1,0,1,0	43/13860
a		(2,5),2(3,8)≻(8,21)	1/1386
83	4,0,0,1,0,1,0,0,1,0	6,0,0,4,0,5,1,0,5,0,0,1,1,0,0	5/1848
a		(3,11),(1,4)≻(4,15)	1/840
b		(2,5),(3,8)≻(5,13)	47/60060
84	4,0,0,1,1,1,1,0,1,1	8,0,0,4,0,1,3,0,5,0,0,1,0,0,1	5/1848
a		(2,5),(3,8)≻(5,13)	47/60060

85	4,0,0,0,1,0,1,0,0,0	11,0,0,3,0,1,1,1,6,0,1,0,1,1,0	71/27720
a		(3,8),(4,11)≻(7,19)	47/23940
86	4,0,0,1,0,1,0,0,1,0	6,0,0,4,0,5,0,1,4,0,1,0,2,0,0	1/462
87	4,0,0,0,1,0,1,0,1,0	11,0,0,3,0,1,1,1,5,1,0,0,2,0,1	31/9240
a		(2,5),(3,8)≻(5,13)	43/30030
b		(3,8),(4,11)≻(7,19)	11/3990
c		(4,11),(1,3)≻(5,14)	1/840
d		(1,3),(3,10)≻(4,13)	19/24024
88	4,0,0,0,1,0,1,0,1,1	11,0,0,3,0,1,1,1,6,0,0,1,1,0,1	3/616
a		(3,11),(1,4)≻(4,15);(2,5),(3,8)≻(5,13)	43/30030
b		(3,11),(1,4)≻(4,15);(3,8),(4,11)≻(7,19)	11/3990
c		(3,11),(1,4)≻(4,15);(4,11),(1,3)≻(5,14)	1/840
d		(2,5),(3,8)≻(5,13);(4,11),(1,3)≻(5,14)	47/60060
e		(4,11),2(1,3)≻(6,17)	41/31416
89	4,0,0,1,1,1,1,0,1,1	8,0,0,4,0,1,2,1,4,0,1,0,1,0,1	1/462
a		2(3,8),(4,11)≻(10,27)	1/756
90	4,0,0,0,1,0,1,0,1,1	11,0,0,3,0,1,0,2,5,0,1,0,2,0,1	1/231
a		2(4,11),(1,3)≻(9,25)	1/525
b		(4,11),2(1,3)≻(6,17)	1/1309
91	4,0,0,1,1,1,1,1,1,1	8,0,0,4,0,2,0,2,4,0,1,0,0,1,0	4/3465
92	4,0,0,0,1,0,1,0,0,0	10,0,1,2,0,1,2,0,7,0,1,0,2,0,1	1/252
a		(2,5),2(3,8)≻(8,21)	1/630
b		(4,9),2(3,7)≻(10,23)	5/1932
93	4,0,0,0,0,0,1,0,1,0	9,0,1,1,0,5,1,0,6,0,0,1,2,0,0	37/5544
a		(4,9),(3,7)≻(7,16);(3,11),2(1,4)≻(5,19);(2,5),(3,8)≻(5,13)	27/19760
b		(4,9),(3,7)≻(7,16);5(2,5),(3,8)≻(13,33)	1/528
c		5(2,5),(3,8)≻(13,33);(3,11),(1,4)≻(4,15)	19/13860
d		(3,11),2(1,4)≻(5,19),3(2,5),(3,8)≻(9,23)	4/3919
94	4,0,0,0,1,0,1,1,0,0	10,0,1,2,0,2,1,0,8,0,0,1,0,1,0	97/27720
a		2(2,5),(3,8)≻(7,18)	1/1386
b		(4,9),2(3,7)≻(10,23)	12/5663
95	4,0,0,1,0,1,0,1,1,0	5,0,1,3,0,6,0,0,6,0,0,1,1,0,0	43/13860
a		(3,11),(1,4)≻(4,15)	1/630
b		(4,9),3(3,7)≻(13,30)	1/660
96	4,0,0,1,0,1,1,0,1,0	6,0,1,2,0,5,2,0,4,0,1,0,2,0,0	1/252
a		(4,9),2(3,7)≻(10,23) ;(2,5),(3,8)≻(5,13)	21/31583
b		2(2,5),(3,8)≻(7,18)	1/840
c		(2,5),2(3,8)≻(8,21)	1/630
97	4,0,0,1,1,1,1,1,1,1	7,0,1,3,0,2,2,0,6,0,0,1,0,0,1	43/13860
a		(4,9),3(3,7)≻(13,30)	1/660
b		(2,5),2(3,8)≻(8,21)	1/1386
98	4,0,0,0,0,0,1,0,1,0	9,0,1,1,0,5,0,1,5,0,1,0,3,0,0	17/2772
a		(4,9),(3,7)≻(7,16);2(1,3),(4,11)≻(6,17)	3/1904
b		4(1,3),(4,11)≻(8,23)	5/5796
99	4,0,0,0,1,0,1,1,0,0	10,0,1,2,0,2,0,1,7,0,1,0,1,1,0	41/13860
a		(4,9),2(3,7)≻(10,23)	33/20914
b		(1,4),(2,9)≻(3,13)	37/45045
c		(1,3),(4,11)≻(5,14)	1/1260
100	4,0,0,0,1,0,1,1,1,0	10,0,1,2,0,2,0,1,6,1,0,0,2,0,1	13/3465
a		(4,11),2(3,7)≻(10,23)	3/1265
b		(1,3),(3,10)≻(4,13)	27/22733
c		(4,11),(1,3)≻(5,14)	1/630
101	4,0,0,0,1,0,1,1,1,1	10,0,1,2,0,2,0,1,7,0,0,1,1,0,1	73/13860
a		(3,11),(1,4)≻(4,15);(4,9),2(3,7)≻(10,23)	3/1265
b		(3,11),(1,4)≻(4,15); (1,3),(4,11)≻(5,14)	1/630
c		(4,9),2(3,7)≻(10,23); (1,3),(4,11)≻(5,14)	29/16839
d		3(1,3),(4,11)≻(7,20)	1/1386

e		(4,9),(3,7)≻(7,16);(4,11),2(1,3)≻(6,17)	19/26767
102	4,0,0,1,1,1,1,1,1,1	7,0,1,3,0,2,1,1,5,0,1,0,1,0,1	71/27720
a		(4,9),3(3,7)≻(13,30)	3/3080
b		(3,8),(4,11)≻(7,19);(4,9),3(3,7)≻(13,30)	31/31920
103	4,0,0,0,0,0,1,0,1,0	8,1,0,1,0,5,1,0,6,0,1,0,3,0,0	3/616
a		5(2,5),(3,8)≻(13,33)	1/924
b		(1,2),(5,11)≻(6,13)	1/728
104	4,0,0,0,1,0,1,1,0,0	9,1,0,2,0,2,1,0,8,0,1,0,1,1,0	47/27720
105	4,0,0,1,0,1,0,1,1,0	4,1,0,3,0,6,0,0,6,0,1,0,2,0,0	1/770
106	4,0,0,0,1,0,1,1,1,0	9,1,0,2,0,2,1,0,7,1,0,0,2,0,1	23/9240
107	4,0,0,0,1,0,1,1,1,1	9,1,0,2,0,2,1,0,8,0,0,1,1,0,1	37/9240
a		(3,11),(1,4)≻(4,15)	23/9240
b		2(2,5),(3,8)≻(7,18)	17/13860
108	4,0,0,1,1,1,1,1,1,1	6,1,0,3,0,2,2,0,6,0,1,0,1,0,1	1/770
109	4,0,0,0,1,0,1,1,1,1	9,1,0,2,0,2,0,1,7,0,1,0,2,0,1	4/1155
a		(4,11),(1,3)≻(5,14)	1/770
110	4,0,0,0,1,0,1,1,0,0	9,0,2,1,0,2,1,0,8,0,1,0,2,0,1	11/2520
a		2(4,9),(3,7)≻(11,25);(2,5),(3,8)≻(5,13)	8/6825
b		2(2,5),(3,8)≻(7,18)	1/630
111	4,0,0,0,0,0,1,1,1,0	8,0,2,0,0,6,0,0,7,0,0,1,2,0,0	7/990
a		(3,11),2(1,4)≻(5,19)	4/855
112	4,0,0,1,0,1,1,1,1,0	5,0,2,1,0,6,1,0,5,0,1,0,2,0,0	11/2520
a		2(4,9),(3,7)≻(11,25);(2,5),(3,8)≻(5,13)	8/6825
b		4(2,5),(3,8)≻(11,28)	1/1260
113	4,0,0,0,0,0,1,1,1,0	7,1,1,0,0,6,0,0,7,0,1,0,3,0,0	73/13860
a		(5,11),(4,9)≻(9,20)	1/210
b		(1,2),(5,11)≻(6,13)	1/945
114	4,0,0,0,1,0,0,0,1,0	11,0,0,3,0,2,1,0,7,0,1,1,0,0,1	47/9240
a		2(2,5),(3,8)≻(7,18);(2,7),(3,11)≻(5,18)	1/630
115	4,0,0,0,1,0,0,0,1,0	11,0,0,3,0,2,0,1,6,0,2,0,1,0,1	1/220
a		2(1,3),(4,11)≻(6,17)	1/1020
116	4,0,0,0,1,0,1,0,1,0	12,0,0,1,1,1,1,1,5,0,2,0,0,1,0	71/27720
a		(3,7),(5,12)≻(8,19);(3,8),(4,11)≻(7,19)	8/5985
b		(5,12),(2,5)≻(7,17);(3,8),(4,11)≻(7,19)	20/20349
117	4,0,0,0,1,0,1,0,1,0	12,0,0,2,0,2,1,1,5,0,2,0,0,1,0	137/27720
a		2(2,5),(3,8)≻(7,18)	1/462
b		(2,5),(3,8)≻(5,13);(4,11),(1,3)≻(5,14)	1/1170
c		(3,8),(4,11)≻(7,19)	26/5985
d		(4,11),2(1,3)≻(6,17)	59/42840
118	4,0,0,0,1,0,1,0,1,0	11,0,1,0,1,1,2,0,6,0,2,0,1,0,1	1/252
a		(5,12),(2,5)≻(7,17)	16/5355
b		(2,5),2(3,8)≻(8,21)	1/630
119	4,0,0,0,1,0,0,1,1,0	10,0,1,2,0,3,0,0,8,0,1,1,0,0,1	19/3465
a		(4,9),2(3,7)≻(10,23);(2,7),(3,11)≻(5,18)	7/2070
120	4,0,0,0,1,0,1,0,1,0	11,0,1,1,0,2,2,0,6,0,2,0,1,0,1	2/315
a		(4,9),(3,7)≻(7,16);2(2,5),2(3,8)≻2(5,13)	11/7280
121	4,0,0,0,1,1,1,0,1,0	9,0,1,2,0,2,2,0,4,1,1,0,2,0,0	1/252
a		(4,9),2(3,7)≻(10,23);(3,10),(2,7)≻(5,17)	41/23460
b		(4,9),2(3,7)≻(10,23);(2,5),(3,8)≻(5,13)	21/31583
c		(4,9),(3,7)≻(7,16);(3,10),(2,7)≻(5,17)	61/28560
d		2(2,5),(3,8)≻(7,18)	1/840
e		(2,5),2(3,8)≻(8,21);(3,10),(2,7)≻(5,17)	4/5355
f		(1,3),(3,10)≻(4,13)	23/16380
122	4,0,0,0,1,1,1,0,1,1	9,0,1,2,0,2,2,0,5,0,1,1,1,0,0	19/3465
a		(4,9),2(3,7)≻(10,23);(2,7),(3,11)≻(5,18);(2,5),2(3,8)≻(8,21)	29/28980
b		(4,9),2(3,7)≻(10,23);(3,11),(1,4)≻(4,15);(2,5),(3,8)≻(5,13)	21/31583
c		(4,9),(3,7)≻(7,16);,2(2,5),2(3,8)≻2(5,13)	31/48048

d		(4,9),(3,7)≻(7,16);(2,7),(3,11)≻(5,18),2(2,5),(3,8)≻(7,18)	1/1008
e		(4,9),(3,7)≻(7,16); (3,11),(1,4)≻(4,15); (2,5),(3,8)≻(5,13)	23/21840
e		(2,7),(3,11)≻(5,18);2(2,5),2(3,8)≻2(5,13)	1/1092
123	4,0,0,0,1,0,1,1,1,0	11,0,1,0,1,2,0,1,6,0,2,0,0,1,0	41/13860
a		(5,12),2(2,5)≻(9,22)	1/693
b		(4,11),(1,3)≻(5,14)	1/1260
124	4,0,0,0,1,0,1,1,1,0	11,0,1,1,0,3,0,1,6,0,2,0,0,1,0	37/6930
a		(4,9),(3,7)≻(7,16);2(1,3),(4,11)≻(6,17)	31/39643
b		3(1,3),(4,11)≻(7,20)	31/39643
125	4,0,0,0,1,1,1,0,1,1	9,0,1,2,0,2,1,1,4,0,2,0,2,0,0	137/27720
a		(4,9),2(3,7)≻(10,23);2(2,5),(3,8)≻(7,18)	25/31878
b		(4,9),2(3,7)≻(10,23);(4,11),(1,3)≻(5,14)	9/6440
c		(4,9),2(3,7)≻(10,23);(3,8),(4,11)≻(7,19)	59/19906
d		(4,11),2(1,3)≻(6,17)	59/42840
e		2(2,5),(3,8)≻(7,18);(4,11),(1,3)≻(5,14)	1/1170
f		(4,9),(3,7)≻(7,16);(4,11),(1,3)≻(5,14)	1/560
126	4,0,0,0,1,0,0,1,1,0	9,1,0,2,0,3,0,0,8,0,2,0,1,0,1	17/4620
127	4,0,0,0,1,0,1,1,1,0	10,1,0,0,1,2,1,0,7,0,2,0,0,1,0	47/27720
a		(5,12),(2,5)≻(7,17)	68/95087
128	4,0,0,0,1,0,1,1,1,0	10,1,0,1,0,3,1,0,7,0,2,0,0,1,0	42/10303
a		3(2,5),(3,8)≻(9,23)	13/15939
129	4,0,0,0,1,1,1,0,1,1	8,1,0,2,0,2,2,0,5,0,2,0,2,0,0	17/4620
a		2(2,5),(3,8)≻(7,18)	5/5544
b		(2,5),2(3,8)≻(8,21)	1/770
130	4,0,0,0,1,0,1,1,1,0	10,0,2,0,0,3,1,0,7,0,2,0,1,0,1	17/2520
a		3(2,5),(3,8)≻(9,23)	101/28980
131	4,0,0,0,1,1,1,1,1,0	8,0,2,1,0,3,1,0,5,1,1,0,2,0,0	11/2520
a		2(4,9),(3,7)≻(11,25);(2,5),(3,8)≻(5,13)	13/4200
b		2(4,9),(3,7)≻(11,25);(3,10),(2,7)≻(5,17)	23/10200
c		(4,9),(3,7)≻(7,16);(1,3),(3,10)≻(4,13)	31/38323
d		3(2,5),(3,8)≻(9,23)	8/7245
e		2(2,5),(3,8)≻(7,18),(3,10),(2,7)≻(5,17)	4/5355
f		(1,3),(3,10)≻(4,13)	59/32760
132	4,0,0,0,1,1,1,1,1,1	8,0,2,1,0,3,1,0,6,0,1,1,1,0,0	49/8333
a		2(4,9),(3,7)≻(11,25);(2,7),(3,11)≻(5,18);2(2,5),(3,8)≻(7,18)	1/900
b		2(4,9),(3,7)≻(11,25);3(2,5),(3,8)≻(9,23)	239/177100
c		(4,9),(3,7)≻(7,16);(2,7),(3,11)≻(5,18);3(2,5),(3,8)≻(9,23)	1/1104
d		2(4,9),(3,7)≻(11,25);(3,11),(1,4)≻(4,15);(2,5),(3,8)≻(5,13)	8/6825
e		(3,11),(1,4)≻(4,15); 3(2,5),(3,8)≻(9,23)	8/7245
133	4,0,0,0,1,1,1,1,1,1	8,0,2,1,0,3,0,1,5,0,2,0,2,0,0	37/6930
a		2(4,9),(3,7)≻(11,25);(4,11),(1,3)≻(5,14)	1/525
b		(4,9),(3,7)≻(7,16); (4,11),2(1,3)≻(6,17)	31/39643
c		(4,11),3(1,3)≻(7,20)	1/1260
134	4,0,0,0,1,1,1,1,1,1	7,1,1,1,0,3,1,0,6,0,2,0,2,0,0	42/10303
a		(5,11),(4,9)≻(9,20);2(2,5),(3,8)≻(7,18)	1/1260
b		(4,9),(3,7)≻(7,16); (2,5),(3,8)≻(5,13)	40/34443
c		3(2,5),(3,8)≻(9,23)	13/15939
135	4,0,0,0,1,1,0,0,1,0	9,0,1,2,0,3,1,0,5,0,3,0,1,0,0	13/2520
a		(4,9),2(3,7)≻(10,23);2(2,5),(3,8)≻(7,18)	29/28980
b		(4,9),(3,7)≻(7,16);3(2,5),(3,8)≻(9,23)	1/1104
136	4,0,0,0,1,1,0,1,1,0	8,0,2,1,0,4,0,0,6,0,3,0,1,0,0	1/180
a		2(4,9),(3,7)≻(11,25)	3/700
137	5,0,0,0,1,0,1,0,1,0	14,0,0,3,0,3,1,1,7,0,1,1,0,1,0	42/10303
a		(2,7),(3,11)≻(5,18);(2,5),(3,8)≻(5,13)	43/30030
b		(2,7),(3,11)≻(5,18);(4,11),(1,3)≻(5,14)	1/840
c		(2,7),(3,11)≻(5,18);(3,8),(4,11)≻(7,19)	11/3990

d		3(2,5),(3,8)≻(9,23)	13/15939
e		(4,11),(1,3)≻(5,14)	53/27720
138	5,0,0,0,1,0,1,0,1,0	14,0,0,3,0,3,0,2,6,0,2,0,1,1,0	7/1980
a		(1,4),(2,9)≻(3,13)	1/715
b		2(4,11),(1,3)≻(9,25)	1/900
139	5,0,0,0,1,0,1,0,1,0	13,0,1,2,0,3,2,0,8,0,1,1,1,0,1	19/3465
a		(4,9),2(3,7)≻(10,23);(2,7),(3,11)≻(5,18);(2,5),2(3,8)≻(8,21)	29/28980
b		(4,9),2(3,7)≻(10,23);(3,11),(1,4)≻(4,15);(2,5),(3,8)≻(513)	21/31583
c		(4,9),(3,7)≻(7,16);(2,7),(3,11)≻(5,18);2(2,5),(3,8)≻(7,18)	1/1008
d		(3,11),(1,4)≻(4,15); (2,5),2(3,8)≻(8,21)	1/630
e		(3,11),(1,4)≻(4,15);3(2,5),(3,8)≻(7,18)	41/57960
f		(2,7),(3,11)≻(5,18);3(2,5),(3,8)≻(9,23)	29/19320
g		(2,7),(3,11)≻(5,18);2(2,5),2(3,8)≻2(5,13)	1/1092
140	5,0,0,0,1,0,1,0,1,0	13,0,1,2,0,3,1,1,7,0,2,0,2,0,1	137/27720
a		(4,9),2(3,7)≻(10,23);2(2,5),(3,8)≻(7,18)	25/31878
b		(4,9),2(3,7)≻(10,23);(4,11),(1,3)≻(5,14)	9/6440
c		(4,9),2(3,7)≻(10,23);(3,8),(4,11)≻(7,19)	59/19906
d		(4,9),(3,7)≻(7,16);3(2,5),(3,8)≻(9,23)	17/24661
e		(2,5),(3,8)≻(5,13);(4,11),(1,3)≻(5,14)	1/1170
f		2(1,3),(4,11)≻(6,17)	59/42840
141	5,0,0,0,1,0,1,1,1,0	13,0,1,2,0,4,0,1,8,0,1,1,0,1,0	31/6930
a		(4,9),2(3,7)≻(10,23);(2,7),(3,11)≻(5,18)	3/1808
b		(4,9),2(3,7)≻(10,23);(1,3),(4,11)≻(5,14)	25/26924
c		(2,7),(3,11)≻(5,18);(1,3),(4,11)≻(5,14)	1/630
142	5,0,0,0,1,0,1,0,1,0	12,1,0,2,0,3,2,0,8,0,2,0,2,0,1	17/4620
a		2(2,5),(3,8)≻(7,18)	5/5544
b		(2,5),2(3,8)≻(8,21)	1/770
143	5,0,0,0,1,0,1,1,1,0	12,1,0,2,0,4,1,0,9,0,1,1,0,1,0	89/27720
a		(2,7),(3,11)≻(5,18)	23/9240
b		(2,5),(3,8)≻(5,13)	58/45045
144	5,0,0,0,1,0,1,1,1,0	12,1,0,2,0,4,0,1,8,0,2,0,1,1,0	37/13860
145	5,0,0,0,1,0,1,1,1,0	12,0,2,1,0,4,1,0,9,0,1,1,1,0,1	49/8333
a		(2,7),(3,11)≻(5,18);2(4,9),(3,7)≻(11,25);2(2,5),(3,8)≻(7,18)	1/900
b		(2,7),(3,11)≻(5,18);(4,9),(3,7)≻(7,16);3(2,5),(3,8)≻(9,23)	1/1104
c		(2,7),(3,11)≻(5,18); 4(2,5),(3,8)≻(11,28)	1/630
d		(3,11),(1,4)≻(4,15);2(4,9),(3,7)≻(11,25);(2,5),(3,8)≻(5,13)	8/6825
e		(3,11),(1,4)≻(4,15); 4(2,5),(3,8)≻(11,28)	1/1260
f		(3,11),(1,4)≻(4,15);(4,9),(3,7)≻(7,16);(2,5),(3,8)≻(5,13)	19/13104
g		2(4,9),(3,7)≻(11,25); 4(2,5),(3,8)≻(11,28)	2/1925
146	5,0,0,0,1,0,1,1,1,0	12,0,2,1,0,4,0,1,8,0,2,0,2,0,1	37/6930
a		2(4,9),(3,7)≻(11,25);(4,11),(1,3)≻(5,14)	1/525
b		(4,9),(3,7)≻(7,16);(4,11),2(1,3)≻(6,17)	31/39643
c		(4,11),3(1,3)≻(7,20)	1/1260
147	5,0,0,0,1,0,1,1,1,0	11,1,1,1,0,4,1,0,9,0,2,0,2,0,1	42/10303
a		(5,11),(4,9)≻(9,20);2(2,5),(3,8)≻(7,18)	1/1260
b		(4,9),(3,7)≻(7,16);(2,5),(3,8)≻(5,13)	19/6160
c		3(2,5),(3,8)≻(9,23)	13/15939
148	5,0,0,0,1,1,1,0,1,0	11,0,2,1,0,4,2,0,6,0,3,0,2,0,0	1/180
a		2(4,9),(3,7)≻(11,25);4(2,5),(3,8)≻(11,28)	1/1400
b		2(4,9),(3,7)≻(11,25);(2,5),2(3,8)≻(8,21)	1/525
c		3(2,5),2(3,8)≻(12,31)	1/1395
149	5,0,0,0,1,1,1,1,1,0	10,0,3,0,0,5,1,0,7,0,3,0,2,0,0	1/168
a		5(2,5),(3,8)≻(13,33)	1/462

3.3. **Classification of Case 2.** Compared to Case (i), the difference is that we have $\epsilon = 2$ and $\sigma_5 = 1$. So we may do parallel calculations. Since the expressions for $B^{(n)}$ are different from before, we should still list them here in order to justify our result.

$$B^{(0)}\begin{cases} n^0_{1,2} = 5\chi - 4P_3 + P_4 \\ n^0_{1,3} = 4\chi + 2P_3 - 3P_4 + P_5 \\ n^0_{1,4} = \chi + P_3 + 2P_4 - P_5 - 1 \\ n^0_{1,5} = 0 \\ n^0_{1,6} = 1 \end{cases}$$

$$\epsilon_5 = \Delta^5(B^{(0)}) - \Delta^5(B) = 2\chi - P_3 + 2P_5 - P_6 - 1$$

$$B^{(5)}\begin{cases} n^5_{1,2} = 3\chi - 3P_3 + P_4 - 2P_5 + P_6 + 1 \\ n^5_{2,5} = 2\chi - P_3 + 2P_5 - P_6 - 1 \\ n^5_{1,3} = 2\chi + 3P_3 - 3P_4 - P_5 + P_6 + 1 \\ n^5_{1,4} = \chi + P_3 + 2P_4 - P_5 - 1 \\ n^5_{1,5} = 0 \\ n^5_{1,6} = 1 \end{cases}$$

$$\epsilon_7 = \chi - P_3 + P_6 + P_7 - P_8 - 1$$

$$B^{(7)}\begin{cases} n^7_{1,2} = 2\chi - 2P_3 + P_4 - 2P_5 - P_7 + P_8 + 2 + \eta \\ n^7_{3,7} = \chi - P_3 + P_6 + P_7 - P_8 - 1 - \eta \\ n^7_{2,5} = \chi + 2P_5 - 2P_6 - P_7 + P_8 + \eta \\ n^7_{1,3} = 2\chi + 3P_3 - 3P_4 - P_5 + P_6 + 1 - \eta \\ n^7_{2,7} = \eta \\ n^7_{1,4} = \chi + P_3 + 2P_4 - P_5 - 1 - \eta \\ n^7_{1,5} = 0 \\ n^7_{1,6} = 1 \end{cases}$$

$$\epsilon_8 = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 - 1$$

$$B^{(8)}\begin{cases} n^8_{1,2} = 2\chi - 2P_3 + P_4 - 2P_5 - P_7 + P_8 + 2 + \eta \\ n^8_{3,7} = \chi - P_3 + P_6 + P_7 - P_8 - 1 - \eta \\ n^8_{2,5} = \chi + P_3 + P_4 + P_5 - 3P_6 - P_7 + P_9 + 1 + \eta \\ n^8_{3,8} = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 - 1 \\ n^8_{1,3} = 2\chi + 4P_3 - 2P_4 - 2P_5 - P_8 + P_9 + 2 - \eta \\ n^8_{2,7} = \eta \\ n^8_{1,4} = \chi + P_3 + 2P_4 - P_5 - 1 - \eta \\ n^8_{1,5} = 0 \\ n^8_{1,6} = 1 \end{cases}$$

$$\epsilon_9 = -2P_3 + P_4 + P_5 - P_7 + P_8 + P_9 - P_{10} - 1 + \eta$$

$$B^{(9)}\begin{cases} n^9_{1,2} = 2\chi - 3P_5 - P_9 + P_{10} + 3 \\ n^9_{4,9} = -2P_3 + P_4 + P_5 - P_7 + P_8 + P_9 - P_{10} - 1 + \eta \\ n^9_{3,7} = \chi + P_3 - P_4 - P_5 + P_6 + 2P_7 - 2P_8 - P_9 + P_{10} - 2\eta \\ n^9_{2,5} = \chi + P_3 + P_4 + P_5 - 3P_6 - P_7 + P_9 + 1 + \eta \\ n^9_{3,8} = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 - 1 \\ n^9_{1,3} = 2\chi + 4P_3 - 2P_4 - 2P_5 - P_8 + P_9 + 2 - \eta \\ n^9_{2,7} = \eta \\ n^9_{1,4} = \chi + P_3 + 2P_4 - P_5 - 1 - \eta \\ n^9_{1,5} = 0 \\ n^9_{1,6} = 1 \end{cases}$$

$$\epsilon_{10} = -P_3 + 2P_6 + P_{10} - P_{11} - 2 - \eta$$

$$B^{(10)}\begin{cases} n^{10}_{1,2} = 2\chi - 3P_5 - P_9 + P_{10} + 3 \\ n^{10}_{4,9} = -2P_3 + P_4 + P_5 - P_7 + P_8 + P_9 - P_{10} - 1 + \eta \\ n^{10}_{3,7} = \chi + P_3 - P_4 - P_5 + P_6 + 2P_7 - 2P_8 - P_9 + P_{10} - 2\eta \\ n^{10}_{2,5} = \chi + P_3 + P_4 + P_5 - 3P_6 - P_7 + P_9 + 1 + \eta \\ n^{10}_{3,8} = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 - 1 \\ n^{10}_{1,3} = 2\chi + 5P_3 - 2P_4 - 2P_5 - 2P_6 - P_8 + P_9 - P_{10} + P_{11} + 4 \\ n^{10}_{3,10} = -P_3 + 2P_6 + P_{10} - P_{11} - 2 - \eta \\ n^{10}_{2,7} = P_3 - 2P_6 - P_{10} + P_{11} + 2 + 2\eta \\ n^{10}_{1,4} = \chi + P_3 + 2P_4 - P_5 - 1 - \eta \\ n^{10}_{1,5} = 0 \\ n^{10}_{1,6} = 1 \end{cases}$$

By computing $\Delta^{11}(B^{(10)})$, We get

$$\epsilon_{11} = \chi + P_3 - P_5 - P_8 + P_{10} + P_{11} - 2 - \eta$$

$$B^{(11)}\begin{cases} n^{11}_{1,2} = 2\chi - 3P_5 - P_9 + P_{10} + 3 - \alpha \\ n^{11}_{5,11} = \alpha \\ n^{11}_{4,9} = -2P_3 + P_4 + P_5 - P_7 + P_8 + P_9 - P_{10} - 1 + \eta - \alpha \\ n^{11}_{3,7} = \chi + P_3 - P_4 - P_5 + P_6 + 2P_7 - 2P_8 - P_9 + P_{10} - 2\eta \\ n^{11}_{2,5} = \chi + P_3 + P_4 + P_5 - 3P_6 - P_7 + P_9 + 1 + \eta \\ n^{11}_{3,8} = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 - 1 - \beta \\ n^{11}_{4,11} = \beta \\ n^{11}_{1,3} = 2\chi + 5P_3 - 2P_4 - 2P_5 - 2P_6 - P_8 + P_9 - P_{10} + P_{11} + 4 - \beta \\ n^{11}_{3,10} = -P_3 + 2P_6 + P_{10} - P_{11} - 2 - \eta \\ n^{11}_{2,7} = -\chi + P_5 - 2P_6 + P_8 - 2P_{10} + 4 + 3\eta + \alpha + \beta \\ n^{11}_{3,11} = \chi + P_3 - P_5 - P_8 + P_{10} + P_{11} - 2 - \eta - \alpha - \beta \\ n^{11}_{1,4} = 2P_4 + P_8 - P_{10} - P_{11} + 1 + \alpha + \beta \\ n^{11}_{1,5} = 0 \\ n^{11}_{1,6} = 1 \end{cases}$$

$$\epsilon_{12} = -\chi - 3P_3 + 2P_5 + P_6 - P_7 + P_8 - P_{13} - 1 - \eta$$

$$B^{(12)}\begin{cases} n^{12}_{1,2} = 2\chi - 3P_5 - P_9 + P_{10} + 3 - \alpha \\ n^{12}_{5,11} = \alpha \\ n^{12}_{4,9} = -2P_3 + P_4 + P_5 - P_7 + P_8 + P_9 - P_{10} - 1 + \eta - \alpha \\ n^{12}_{3,7} = 2\chi + 4P_3 - P_4 - 3P_5 + 3P_7 - 3P_8 - P_9 + P_{10} + P_{13} + 1 - \eta \\ n^{12}_{5,12} = -\chi - 3P_3 + 2P_5 + P_6 - P_7 + P_8 - P_{13} - 1 - \eta \\ n^{12}_{2,5} = 2\chi + 4P_3 + P_4 - P_5 - 4P_6 - P_8 + P_9 + P_{13} + 2 + 2\eta \\ n^{12}_{3,8} = -P_3 - P_4 + P_5 + P_6 + P_8 - P_9 - 1 - \beta \\ n^{12}_{4,11} = \beta \\ n^{12}_{1,3} = 2\chi + 5P_3 - 2P_4 - 2P_5 - 2P_6 - P_8 + P_9 - P_{10} + P_{11} + 4 - \beta \\ n^{12}_{3,10} = -P_3 + 2P_6 + P_{10} - P_{11} - 2 - \eta \\ n^{12}_{2,7} = -\chi + P_5 - 2P_6 + P_8 - 2P_{10} + 4 + 3\eta + \alpha + \beta \\ n^{12}_{3,11} = \chi + P_3 - P_5 - P_8 + P_{10} + P_{11} - 2 - \eta - \alpha - \beta \\ n^{12}_{1,4} = 2P_4 + P_8 - P_{10} - P_{11} + 1 + \alpha + \beta \\ n^{12}_{1,5} = 0 \\ n^{12}_{1,6} = 1 \end{cases}$$

Similarly we have the boundedness for χ, η, α, β and P_{13}, therefore the boundedness of $B^{(12)}$. Here we have:

$$\chi \leq 3. \tag{3.9}$$

$$\eta \leq \chi + 2. \tag{3.10}$$

$$\alpha \leq 3 + \eta. \tag{3.11}$$

$$\beta \leq 2 \tag{3.12}$$

Finally, we have

$$P_{13} \leq -\chi + 4. \tag{3.13}$$

The result after running our computer program is unexpectedly simple.

No	$(\chi, P_3, P_4, \ldots, P_{11})$	$B^{(12)} = (n^{12}_{1,2}, n^{12}_{5,11}, \ldots, n^{12}_{1,4}, n^{12}_{1,6})$	K^3
1	2,0,0,1,1,0,1,0,1,1	5,0,0,1,0,1,2,0,3,0,0,0,0,1	1/420
2	2,0,0,1,1,0,1,1,1,1	4,0,1,0,0,2,1,0,4,0,0,0,0,1	1/360
a		(2,5),(3,8)$\succ$(5,13)	1/1170

Theorem 3.5. *Let X be a nonsingular projective 3-fold of general type with the pluricanonical section index $\delta(X) = 12$. Then* $\mathrm{Vol}(X) \geq \frac{31}{48048}$ *and the equality holds if and only it corresponds to Case 122.c.*

Proof. When $P_{12} \geq 3$, one has $Vol(X) \geq \frac{1}{1224}$ by those inequalities in Chen–Chen [3, Section 3].

If $P_{12} = 2$, we can get that $Vol(X) \geq \frac{31}{48048}$ from our tables and the equality holds if and only if it corresponds to Case 122.c. □

Remark 3.6. It is mysterious whether the lower bound $\frac{31}{48048}$ is optimal or not.

References

[1] J. A. Chen, M. Chen, *Explicit birational geometry of threefolds of general type, I.* Ann. Sci. Ec. Norm. Super **43**(2010), 365–394.

[2] J. A. Chen, M. Chen, *Explicit birational geometry of threefolds of general type, II.* J. of Diff. Geom. **86**(2010), 237–271.

[3] J. A. Chen, M. Chen, *Explicit birational geometry of 3-folds and 4-folds of general type, III.* ArXiv: 1302.0374 (v3, 44 pages)

[4] C. D. Hacon, J. McKernan, *Boundedness of pluricanonical maps of varieties of general type*, Invent. Math. 166 (2006), 1–25.

[5] M. Reid, *Young person's guide to canonical singularities*, Proc. Symposia in pure Math. **46**(1987), 345–414.

[6] S. Takayama, *Pluricanonical systems on algebraic varieties of general type*, Invent. Math. 165 (2006), 551–587.

[7] H. Tsuji, *Pluricanonical systems of projective varieties of general type. I.* Osaka J. Math. **43** (2006), no. 4, 967–995.

School of Mathematical Sciences, Fudan University, Shanghai 200433, China
E-mail address: 10300180105@fudan.edu.cn